

2025 Performance and Outcome Measures	JHMC 2025 Q3 Scores	AHA/NY Stroke Goal
Stroke patients who receive prophylaxis for VTE	96%	85%
Patients with an ischemic stroke prescribed antithrombotic therapy at discharge.	100%	85%
Patients with an ischemic stroke with atrial fibrillation/flutter discharged on anticoagulation therapy	100%	85%
Ischemic stroke patients given early antithrombotics	94%	85%
Stroke patients discharged on cholesterol reducing medication	100%	85%
Stroke patients or their caregivers given educational materials during their hospital stay	100%	85%
Stroke patients who were assessed for rehabilitation services.	99%	85%
National Institute of Health Stroke Scale on Admission	95%	85%
Severity measurements performed for subarachnoid hemorrhage and intracranial hemorrhage patients	94%	88%
Rate of rapid effective reperfusion from skin puncture in patients receiving mechanical endovascular reperfusion therapy	89%	79.4%
Aggregated Complication Rate of less than 6% (CEA)	0%	≤6% (TJC)
Aggregated Complication Rate of less than 3% (CEA)	0%	≤3% (TJC)
Aggregated Complication Rate of less than 6% (CAS)	0%	≤6% (TJC)
Aggregated Complication Rate of less than 3% (CAS)	0%	≤3% (TJC)
24-hour Post-procedure stroke and death rate of ≤1% (diagnostic catheter angiography)	0%	≤1% (TJC)